

Diploma Programme

Data booklet



Diploma Programme

Physics
Data booklet

First examinations 2009

International Baccalaureate Organization

Buenos Aires

Cardiff

Geneva

New York

Singapore

Diploma Programme
Physics—data booklet

Published September 2007

International Baccalaureate Organization
Peterson House, Malthouse Avenue, Cardiff Gate
Cardiff, Wales GB CF23 8GL
United Kingdom
Phone: +44 29 2054 7777
Fax: +44 29 2054 7778
Web site: <http://www.ibo.org>

© International Baccalaureate Organization 2007

The International Baccalaureate Organization (IBO) was established in 1968 and is a non-profit, international educational foundation registered in Switzerland.

The IBO is grateful for permission to reproduce and/or translate any copyright material used in this publication. Acknowledgments are included, where appropriate, and, if notified, the IBO will be pleased to rectify any errors or omissions at the earliest opportunity.

IBO merchandise and publications in its official and working languages can be purchased through the IB store at <http://store.ibo.org>. General ordering queries should be directed to the sales and marketing department in Cardiff.

Phone: +44 29 2054 7746
Fax: +44 29 2054 7779
E-mail: sales@ibo.org

Contents

Fundamental constants	1
Metric (SI) multipliers	2
Unit conversions	3
Electrical circuit symbols	4
Equations—Core and AHL	5
Equations—Options SL	11
Equations—Options SL and HL	13
Equations—Options HL	15

Fundamental constants

Quantity	Symbol	Approximate value
Acceleration of free fall (Earth's surface)	g	9.81 m s^{-2}
Gravitational constant	G	$6.67 \times 10^{-11} \text{ N m}^2 \text{ kg}^{-2}$
Avogadro's constant	N_A	$6.02 \times 10^{23} \text{ mol}^{-1}$
Gas constant	R	$8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Boltzmann's constant	k	$1.38 \times 10^{-23} \text{ J K}^{-1}$
Stefan–Boltzmann constant	σ	$5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$
Coulomb constant	k	$8.99 \times 10^9 \text{ N m}^2 \text{ C}^{-2}$
Permittivity of free space	ϵ_0	$8.85 \times 10^{-12} \text{ C}^2 \text{ N}^{-1} \text{ m}^{-2}$
Permeability of free space	μ_0	$4\pi \times 10^{-7} \text{ T m A}^{-1}$
Speed of light in vacuum	c	$3.00 \times 10^8 \text{ m s}^{-1}$
Planck's constant	h	$6.63 \times 10^{-34} \text{ J s}$
Elementary charge	e	$1.60 \times 10^{-19} \text{ C}$
Electron rest mass	m_e	$9.110 \times 10^{-31} \text{ kg} = 0.000549 \text{ u} = 0.511 \text{ MeV c}^{-2}$
Proton rest mass	m_p	$1.673 \times 10^{-27} \text{ kg} = 1.007276 \text{ u} = 938 \text{ MeV c}^{-2}$
Neutron rest mass	m_n	$1.675 \times 10^{-27} \text{ kg} = 1.008665 \text{ u} = 940 \text{ MeV c}^{-2}$
Unified atomic mass unit	u	$1.661 \times 10^{-27} \text{ kg} = 931.5 \text{ MeV c}^{-2}$

Metric (SI) multipliers

Prefix	Abbreviation	Value
tera	T	10^{12}
giga	G	10^9
mega	M	10^6
kilo	k	10^3
hecto	h	10^2
deca	da	10^1
deci	d	10^{-1}
centi	c	10^{-2}
milli	m	10^{-3}
micro	μ	10^{-6}
nano	n	10^{-9}
pico	p	10^{-12}
femto	f	10^{-15}

Unit conversions

$$1 \text{ light year (ly)} = 9.46 \times 10^{15} \text{ m}$$

$$1 \text{ parsec (pc)} = 3.26 \text{ ly}$$
















$$1 \text{ astronomical unit (AU)} = 1.50 \times 10^{11} \text{ m}$$

$$1 \text{ radian (rad)} = \frac{180^\circ}{\pi}$$

$$1 \text{ kilowatt-hour (kWh)} = 3.60 \times 10^6 \text{ J}$$


$$1 \text{ atm} = 1.01 \times 10^5 \text{ N m}^{-2} = 101 \text{ kPa} = 760 \text{ mm Hg}$$

Electrical circuit symbols

cell		battery	
lamp		ac supply	
switch		ammeter	
voltmeter		variable resistor	
resistor		potentiometer	
light-dependent resistor (LDR)		thermistor	
transformer		heating element	
operational amplifier (op-amp)			

Equations—Core and AHL

Note: All equations relate to the magnitude of the quantities only. Vector notation has not been used.

Core	AHL
<p>Topic 1: Physics and physical measurement</p> <p>If $y = a \pm b$ then $\Delta y = \Delta a + \Delta b$</p> <p>If $y = \frac{ab}{c}$ then $\frac{\Delta y}{y} = \frac{\Delta a}{a} + \frac{\Delta b}{b} + \frac{\Delta c}{c}$</p>  <p>$A_H = A \cos \theta$ $A_V = A \sin \theta$</p>	

Core	AHL
<p>Topic 2: Mechanics</p> $s = \frac{u + v}{2}t$ $s = ut + \frac{1}{2}at^2$ $v^2 = u^2 + 2as$ $F = ma$ $p = mv$ $F = \frac{\Delta p}{\Delta t}$ <p>Impulse = $F\Delta t = m\Delta v$</p> $W = Fs \cos \theta$ $E_k = \frac{1}{2}mv^2$ $E_k = \frac{p^2}{2m}$ $\Delta E_p = mg\Delta h$ <p>power = Fv</p> $a = \frac{v^2}{r} = \frac{4\pi^2 r}{T^2}$	
<p>Topic 3: Thermal physics</p> $P = \frac{F}{A}$ $Q = mc\Delta T$ $Q = mL$	<p>Topic 10: Thermal physics</p> $PV = nRT$ $W = P\Delta V$ $Q = \Delta U + W$

Core	AHL
<p>Topic 4: Oscillations and waves</p> $\omega = \frac{2\pi}{T}$ $x = x_0 \sin \omega t; \quad x = x_0 \cos \omega t$ $v = v_0 \cos \omega t; \quad v = -v_0 \sin \omega t$ $v = \pm \omega \sqrt{(x_0^2 - x^2)}$ $E_K = \frac{1}{2} m \omega^2 (x_0^2 - x^2)$ $E_{K(\max)} = \frac{1}{2} m \omega^2 x_0^2$ $E_T = \frac{1}{2} m \omega^2 x_0^2$ $v = f \lambda$ $\frac{n_1}{n_2} = \frac{\sin \theta_2}{\sin \theta_1} = \frac{v_2}{v_1}$ <p>path difference = $n \lambda$</p> <p>path difference = $(n + \frac{1}{2}) \lambda$</p>	<p>Topic 11: Wave phenomena</p> $f' = f \left(\frac{v}{v \pm u_s} \right) \quad \text{moving source}$ $f' = f \left(\frac{v \pm u_o}{v} \right) \quad \text{moving observer}$ $\Delta f = \frac{v}{c} f$ $\theta = \frac{\lambda}{b}$ $\theta = 1.22 \frac{\lambda}{b}$ $I = I_0 \cos^2 \theta$ $n = \tan \phi$

Core	AHL										
<p>Topic 5: Electric currents</p> $K_e = \frac{1}{2}mv^2$ $I = \frac{\Delta q}{\Delta t}$ $R = \frac{V}{I}$ $R = \frac{\rho L}{A}$ $P = VI = I^2R = \frac{V^2}{R}$ $\mathcal{E} = I(R+r)$ $R = R_1 + R_2 + \dots$ $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \dots$	<p>Topic 12: Electromagnetic induction</p> $\Phi = BA \cos \theta$ $\mathcal{E} = Bvl$ $\mathcal{E} = -N \frac{\Delta \Phi}{\Delta t}$ $\frac{I_s}{I_p} = \frac{V_p}{V_s} = \frac{N_p}{N_s}$ $I_{\text{rms}} = \frac{I_0}{\sqrt{2}}$ $V_{\text{rms}} = \frac{V_0}{\sqrt{2}}$ $R = \frac{V_0}{I_0} = \frac{V_{\text{rms}}}{I_{\text{rms}}}$ $P_{\text{max}} = I_0V_0$ $P_{\text{av}} = \frac{1}{2}I_0V_0$										
<p>Topic 6: Fields and forces</p> <table border="1" data-bbox="261 1268 786 1545"> <tbody> <tr> <td data-bbox="261 1268 522 1402">$F = G \frac{m_1m_2}{r^2}$</td> <td data-bbox="522 1268 786 1402">$F = k \frac{q_1q_2}{r^2}$</td> </tr> <tr> <td data-bbox="261 1402 522 1545">$g = \frac{F}{m}$</td> <td data-bbox="522 1402 786 1545">$E = \frac{F}{q}$</td> </tr> </tbody> </table> $F = \frac{q_1q_2}{4\pi\epsilon_0r^2}$ $F = qvB \sin \theta$ $F = BIL \sin \theta$	$F = G \frac{m_1m_2}{r^2}$	$F = k \frac{q_1q_2}{r^2}$	$g = \frac{F}{m}$	$E = \frac{F}{q}$	<p>Topic 9: Motion in fields</p> <table border="1" data-bbox="873 1268 1305 1722"> <tbody> <tr> <td data-bbox="873 1268 1089 1417">$\Delta V = \frac{\Delta E_p}{m}$</td> <td data-bbox="1089 1268 1305 1417">$\Delta V = \frac{\Delta E_p}{q}$</td> </tr> <tr> <td data-bbox="873 1417 1089 1566">$V = -\frac{Gm}{r}$</td> <td data-bbox="1089 1417 1305 1566">$V = \frac{kq}{r} = \frac{q}{4\pi\epsilon_0r}$</td> </tr> <tr> <td data-bbox="873 1566 1089 1722">$g = -\frac{\Delta V}{\Delta r}$</td> <td data-bbox="1089 1566 1305 1722">$E = -\frac{\Delta V}{\Delta x}$</td> </tr> </tbody> </table>	$\Delta V = \frac{\Delta E_p}{m}$	$\Delta V = \frac{\Delta E_p}{q}$	$V = -\frac{Gm}{r}$	$V = \frac{kq}{r} = \frac{q}{4\pi\epsilon_0r}$	$g = -\frac{\Delta V}{\Delta r}$	$E = -\frac{\Delta V}{\Delta x}$
$F = G \frac{m_1m_2}{r^2}$	$F = k \frac{q_1q_2}{r^2}$										
$g = \frac{F}{m}$	$E = \frac{F}{q}$										
$\Delta V = \frac{\Delta E_p}{m}$	$\Delta V = \frac{\Delta E_p}{q}$										
$V = -\frac{Gm}{r}$	$V = \frac{kq}{r} = \frac{q}{4\pi\epsilon_0r}$										
$g = -\frac{\Delta V}{\Delta r}$	$E = -\frac{\Delta V}{\Delta x}$										

Core	AHL
<p data-bbox="272 296 737 323">Topic 7: Atomic and nuclear physics</p> $E = mc^2$	<p data-bbox="829 296 1338 359">Topic 13: Quantum physics and nuclear physics</p> $E = hf$ $hf = \phi + E_{\max}$ $hf = hf_0 + eV$ $p = \frac{h}{\lambda}$ $E_{\text{K}} = \frac{n^2 h^2}{8m_e L^2}$ $\Delta x \Delta p \geq \frac{h}{4\pi}$ $\Delta E \Delta t \geq \frac{h}{4\pi}$ $N = N_0 e^{-\lambda t}$ $A = -\frac{\Delta N}{\Delta t}$ $A = \lambda N = \lambda N_0 e^{-\lambda t}$ $T_{\frac{1}{2}} = \frac{\ln 2}{\lambda}$

Core	AHL
<p data-bbox="282 323 740 386">Topic 8: Energy, power and climate change</p> <p data-bbox="282 441 477 483">power = $\frac{1}{2} A \rho v^3$</p> <p data-bbox="282 491 675 533">power per unit length = $\frac{1}{2} A^2 \rho g v$</p> <p data-bbox="282 558 412 621">$I = \frac{\text{power}}{A}$</p> <p data-bbox="282 646 649 722">albedo = $\frac{\text{total scattered power}}{\text{total incident power}}$</p> <p data-bbox="282 747 412 810">$C_s = \frac{Q}{A \Delta T}$</p> <p data-bbox="282 835 461 877">power = $\sigma A T^4$</p> <p data-bbox="282 894 477 936">power = $e \sigma A T^4$</p> <p data-bbox="282 961 509 1037">$\Delta T = \frac{(I_{\text{in}} - I_{\text{out}}) \Delta t}{C_s}$</p>	

Equations—Options SL

Option A: Sight and wave phenomena

$$f' = f \left(\frac{v}{v \pm u_s} \right) \quad \text{moving source} \quad \theta = \frac{\lambda}{b}$$

$$f' = f \left(\frac{v \pm u_o}{v} \right) \quad \text{moving observer} \quad \theta = 1.22 \frac{\lambda}{b}$$

$$\Delta f = \frac{v}{c} f \quad I = I_0 \cos^2 \theta$$

$$n = \tan \phi$$

Option B: Quantum physics and nuclear physics

$$E = hf \quad \Delta E \Delta t \geq \frac{h}{4\pi}$$

$$hf = \phi + E_{\max} \quad N = N_0 e^{-\lambda t}$$

$$hf = hf_0 + eV \quad A = -\frac{\Delta N}{\Delta t}$$

$$p = \frac{h}{\lambda} \quad A = \lambda N = \lambda N_0 e^{-\lambda t}$$

$$E_k = \frac{n^2 h^2}{8m_e L^2} \quad T_{\frac{1}{2}} = \frac{\ln 2}{\lambda}$$

$$\Delta x \Delta p \geq \frac{h}{4\pi}$$

Option C: Digital technology

$$G = -\frac{R_F}{R}$$

$$G = 1 + \frac{R_F}{R}$$

Option D: Relativity and particle physics

$$\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$\Delta t = \gamma \Delta t_0$$

$$L = \frac{L_0}{\gamma}$$

$$\Delta E \Delta t \geq \frac{h}{4\pi}$$

$$R \approx \frac{h}{4\pi mc}$$

$$E = hf$$

Equations—Options SL and HL

Core (SL and HL)	Extension (HL only)
<p>Option E: Astrophysics</p> $L = \sigma AT^4$ $\lambda_{\max} \text{ (metres)} = \frac{2.90 \times 10^{-3}}{T \text{ (kelvin)}}$ $d \text{ (parsec)} = \frac{1}{p \text{ (arc-second)}}$ $b = \frac{L}{4\pi d^2}$ $m - M = 5 \lg \left(\frac{d}{10} \right)$	$L \propto m^n \quad \text{where } 3 < n < 4$ $\frac{\Delta \lambda}{\lambda} \cong \frac{v}{c}$ $v = H_0 d$
<p>Option F: Communications</p> $n = \frac{1}{\sin C}$ $\text{attenuation / dB} = 10 \lg \frac{I_1}{I_2}$	$G = -\frac{R_F}{R}$ $G = 1 + \frac{R_F}{R}$

Core (SL and HL)	Extension (HL only)
<p>Option G: Electromagnetic waves</p> $\frac{1}{f} = \frac{1}{v} + \frac{1}{u}$ $P = \frac{1}{f}$ $m = \frac{h_i}{h_o} = -\frac{v}{u} \quad M = \frac{\theta_i}{\theta_o}$ $M = \frac{f_o}{f_e}$ $m = \frac{D}{f} + 1 \quad m = \frac{D}{f}$ $s = \frac{\lambda D}{d}$ $\sin \theta = \frac{n\lambda}{d}$ $\frac{x}{D} = \frac{n\lambda}{d}$ $\frac{x}{D} = (n + \frac{1}{2}) \frac{\lambda}{d}$ $d \sin \theta = n\lambda$	$\lambda_{\min} = \frac{hc}{eV}$ $2d \sin \theta = n\lambda$ $2nt = m\lambda$ $2nt = (m + \frac{1}{2})\lambda$ $2nt \cos \phi = m\lambda$ $2nt \cos \phi = (m + \frac{1}{2})\lambda$

Equations—Options HL

Option H: Relativity

$$\gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$$\Delta t = \gamma \Delta t_0$$

$$L = \frac{L_0}{\gamma}$$

$$u'_x = \frac{u_x - v}{1 - \frac{u_x v}{c^2}}$$

$$E_0 = m_0 c^2$$

$$E = \gamma m_0 c^2$$

$$p = \gamma m_0 u$$

$$E_K = (\gamma - 1) m_0 c^2$$

$$E^2 = p^2 c^2 + m_0^2 c^4$$

$$\frac{\Delta f}{f} = \frac{g \Delta h}{c^2}$$

$$R_s = \frac{2GM}{c^2}$$

$$\Delta t = \frac{\Delta t_0}{\sqrt{1 - \frac{R_s}{r}}}$$

Option I: Medical physics

$$IL = 10 \lg \frac{I}{I_0} \quad \text{where} \quad I_0 = 1.0 \times 10^{-12} \text{ W m}^{-2}$$

$$I = I_0 e^{-\mu x}$$

$$\mu x_{\frac{1}{2}} = \ln 2$$

dose equivalent = absorbed dose \times quality factor

$$Z = \rho c$$

$$\frac{1}{T_E} = \frac{1}{T_P} + \frac{1}{T_B}$$

Option J: Particle physics

$$\Delta E \Delta t \geq \frac{h}{4\pi}$$

$$R \approx \frac{h}{4\pi mc}$$

$$E = hf$$

$$E = mc^2 + E_K$$

$$E_a^2 = 2Mc^2 E + (Mc^2)^2 + (mc^2)^2$$

$$\lambda = \frac{h}{p}$$

$$E_K = \frac{3}{2} kT$$